


AMENDMENTS TO THE CLAIMS

Please amend the claims as follows:

1. (Previously Presented) A method of inhibiting human stearyl-CoA desaturase (hSCD) activity comprising contacting a source of hSCD with a compound of formula (I):



wherein:

x and y are each independently 1;

W is -C(O)N(R¹)- or -N(R¹)C(O)-;

V is -C(O)-;

each R¹ is independently selected from the group consisting of hydrogen, C₁-C₁₂alkyl, C₂-C₁₂hydroxyalkyl, C₄-C₁₂cycloalkylalkyl and C₇-C₁₉aralkyl;

R² is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₂-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, aryl, C₇-C₁₉aralkyl, C₃-C₁₂heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl, and C₃-C₁₂heteroarylalkyl;

R³ is phenyl or naphthyl;

R⁴, R⁵ and R⁶ are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or -N(R¹³)₂;


R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰ and R^{10a} are each independently selected from hydrogen or C₁-C₃alkyl;

and

each R¹³ is independently selected from hydrogen or C₁-C₆alkyl;

a stereoisomer, enantiomer or tautomer thereof, or a pharmaceutically acceptable salt thereof.

2. (Currently Amended) A method of treating a disease or condition mediated by stearyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to the mammal in need thereof a therapeutically effective amount of a compound of formula (I):



wherein:

x and y are each independently 1;

W is -C(O)N(R¹)- or -N(R¹)C(O)-;

V is -C(O)-;

each R¹ is independently selected from the group consisting of hydrogen, C₁-C₁₂alkyl, C₂-C₁₂hydroxyalkyl, C₄-C₁₂cycloalkylalkyl and C₇-C₁₉aralkyl;

R² is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₂-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, aryl, C₇-C₁₉aralkyl, C₃-C₁₂heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl, and C₃-C₁₂heteroarylalkyl;

R³ is phenyl or naphthyl;

R⁴, R⁵ and R⁶ are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or -N(R¹³)₂;


R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰ and R^{10a} are each independently selected from hydrogen or C₁-C₃alkyl;

and

each R¹³ is independently selected from hydrogen or C₁-C₆alkyl;
a stereoisomer, enantiomer or tautomer thereof, or a pharmaceutically acceptable salt thereof,

and wherein the disease or condition is selected from the group consisting of Type II diabetes, impaired glucose tolerance, insulin resistance, obesity, fatty liver, non-alcoholic steatohepatitis, dyslipidemia, acne, ~~and metabolic syndrome~~ and any combination of these.

3. (Original) The method of Claim 2 wherein the mammal is a human.
4. (Cancelled).
5. (Currently Amended) The method of Claim ~~[[4]]~~ 3, wherein the disease or condition is Type II diabetes.
6. (Currently Amended) The method of Claim ~~[[4]]~~ 3, wherein the disease or condition is obesity.
7. (Currently Amended) The method of Claim ~~[[4]]~~ 3,4 wherein the disease or condition is ~~metabolic syndrome~~ insulin resistance.
8. (Currently Amended) The method of Claim ~~[[4]]~~ 3, wherein the disease or condition is fatty liver.
9. (Currently Amended) The method of Claim ~~[[4]]~~ 3, wherein the disease or condition is non-alcoholic steatohepatitis.
10. (Currently Amended) A compound of formula (IIa):



wherein:

x and y are each independently 1;

R¹ is selected from the group consisting of hydrogen, C₁-C₁₂alkyl, C₂-C₁₂hydroxyalkyl, C₄-C₁₂cycloalkylalkyl and C₇-C₁₉aralkyl;

R² is selected from the group consisting of C₇-C₁₂alkyl, C₃-C₁₂alkenyl, C₇-C₁₂hydroxyalkyl, C₁-C₁₂alkoxy, C₂-C₁₂alkoxyalkyl, C₃-C₁₂hydroxyalkenyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, C₁₃-C₁₉aralkyl, C₁-C₁₂heteroaryl, C₃-C₁₂heterocyclalkyl and C₃-C₁₂heteroarylalkyl, provided that R² is not pyrazinyl, pyridinonyl, ~~pyrrolidinone~~ pyrrolidinonyl or imidazolyl;

R³ is phenyl or naphthyl;

R⁴, R⁵ and R⁶ are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or -N(R¹³)₂;

R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each independently selected from hydrogen or C₁-C₃alkyl;

and

each R¹³ is independently selected from hydrogen or C₁-C₆alkyl;

a stereoisomer, enantiomer or tautomer thereof, or a pharmaceutically acceptable salt thereof.

11. (Previously Presented) The compound of Claim 10 wherein:

x and y are each 1;

R¹ is hydrogen or C₁-C₆alkyl;

R² is selected from the group consisting of C₇-C₁₂alkyl, C₃-C₁₂alkenyl,

C₇-C₁₂hydroxyalkyl, C₂-C₁₂alkoxyalkyl, C₃-C₁₂hydroxyalkenyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, C₁₃-C₁₉aralkyl, C₃-C₁₂heterocyclalkyl, and C₃-C₁₂heteroarylalkyl;

R³ is phenyl or naphthyl;


R⁴, R⁵ and R⁶ are each hydrogen; and

R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each hydrogen.

12. (Currently Amended) A method of treating a disease or condition mediated by stearoyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 10, and wherein the disease or condition is selected from the group consisting of Type II diabetes, impaired glucose tolerance, insulin resistance, obesity, fatty liver, non-alcoholic steatohepatitis, dyslipidemia, acne, ~~and metabolic syndrome~~ and any combination of these.

13. (Original) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound of Claim 10.

14. (Currently Amended) A compound of formula (IIb):



wherein:

x and y are each independently 1;

R¹ is selected from the group consisting of hydrogen, C₁-C₁₂alkyl, C₂-C₁₂hydroxyalkyl, C₄-C₁₂cycloalkylalkyl and C₇-C₁₉aralkyl;

R² is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl,

C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₁-C₆alkoxy, C₃-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, C₇-C₁₉aralkyl, C₃-C₁₂ heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

or R² is phenyl optionally substituted with one or more substituents selected from halo and C₁-C₆trihaloalkyl;

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆trihaloalkyl, C₁-C₆trihaloalkoxy, C₁-C₆alkylsulfonyl, -N(R¹²)₂, -OC(O)R¹², -C(O)OR¹², -S(O)₂N(R¹²)₂, cycloalkyl, heterocyclyl, heteroaryl and heteroarylcycloalkyl, provided that R³ is not phenyl substituted with optionally substituted thienyl;

R⁴, R⁵ and R⁶ are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or -N(R¹³)₂;

R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each independently selected from hydrogen or C₁-C₃alkyl;

and

each R¹² is independently selected from hydrogen, C₁-C₆alkyl, C₃-C₆cycloalkyl, aryl or aralkyl; and

each R¹³ is independently selected from hydrogen or C₁-C₆alkyl;

a stereoisomer, enantiomer or tautomer thereof, or a pharmaceutically acceptable salt thereof.

15. (Currently Amended) The compound of Claim 14 wherein:

x and y are each 1;

R¹ is hydrogen or C₁-C₆alkyl;

R² is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₁-C₆alkoxy, C₃-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, C₇-C₁₉aralkyl, C₃-C₁₂ heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

or R^2 is phenyl optionally substituted with one or more substituents selected from halo and/or C_1 - C_6 trihaloalkyl;

R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl, C_1 - C_6 trihaloalkoxy, C_1 - C_6 alkylsulfonyl, $-N(R^{12})_2$, $-OC(O)R^{12}$, $-C(O)OR^{12}$ and $-S(O)_2N(R^{12})_2$;

R^4 , R^5 and R^6 are each hydrogen;

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each hydrogen; and

each R^{12} is independently selected from hydrogen, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, aryl or aralkyl.

16. (Original) The compound of Claim 15 wherein:

R^2 is C_7 - C_{12} aralkyl optionally substituted by one or more substituents selected from the group consisting of halo, C_1 - C_3 alkyl and C_1 - C_6 trihaloalkyl; and

R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl and C_1 - C_6 trihaloalkoxy.

17. (Original) The compound of Claim 16 selected from the group consisting of the following:

3-(4-Fluoro-phenyl)-N-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-propionamide;

4-Phenyl-N-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-butyramide;

4-(4-Fluoro-phenyl)-N-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-butyramide; and

3-Phenyl-N-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-propionamide.

18. (Original) The compound of Claim 15 wherein:

R^2 is C_1 - C_{12} alkyl or C_2 - C_{12} alkenyl; and

R^3 is phenyl optionally substituted by one or more substituents selected from the

group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy.

19. (Original) The compound of Claim 18 selected from the group consisting of the following:

Hexanoic acid {5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-amide;

Heptanoic acid {5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-amide; and

5-Methylpentanoic acid {5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-amide.

20. (Original) The compound of Claim 15 wherein:

R² is C₃-C₁₂heteroarylalkyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₃alkyl and C₁-C₆trihaloalkyl; and

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy.

21. (Original) The compound of Claim 20, namely, 3-Pyridin-3-yl-N-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-propionamide.

22. (Original) The compound of Claim 15 wherein:

R² is phenyl optionally substituted with one or more substituents selected from halo and C₁-C₆trihaloalkyl; and

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy.


23. (Original) The compound of Claim 22, namely, 4-Fluoro-N-{5-[4-(2-trifluoromethylbenzoyl)piperazin-1-yl]pyridin-2-yl}benzamide.

24. (Currently Amended) A method of treating a disease or condition mediated by

stearoyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 14, wherein the disease or condition is selected from the group consisting of Type II diabetes, impaired glucose tolerance, insulin resistance, obesity, fatty liver, non-alcoholic steatohepatitis, dyslipidemia, acne, and ~~metabolic syndrome~~ and any combination of these.

25. (Original) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound of Claim 14.

26. (Withdrawn) A compound of formula (III):



wherein:

x and y are each independently 1;

V_a is -C(O)-, -C(S)-, -C(O)N(R¹)-, -C(O)O-, -S(O)₂- or -S(O)₂N(R¹)-;

each R¹ is independently selected from the group consisting of hydrogen, C₁-C₁₂alkyl, C₂-C₁₂hydroxyalkyl, C₄-C₁₂cycloalkylalkyl and C₇-C₁₉aralkyl;

R² is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₁-C₆alkoxy, C₃-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, aryl, C₇-C₁₉aralkyl, C₃-C₁₂ heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

or R² is a multi-ring structure having 2 to 4 rings wherein the rings are independently selected from the group consisting of cycloalkyl, heterocyclyl, aryl and heteroaryl, where some or all of the rings may be fused to each other;

R^3 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_2 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

or R^3 is a multi-ring structure having 2 to 4 rings wherein the rings are independently selected from the group consisting of cycloalkyl, heterocyclyl, aryl and heteroaryl and where some or all of the rings may be fused to each other;

R^4 , R^5 and R^6 are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or $-N(R^{13})_2$;

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each independently selected from hydrogen or C_1 - C_3 alkyl;

and

each R^{13} is independently selected from hydrogen or C_1 - C_6 alkyl;

a stereoisomer, enantiomer or tautomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutical composition thereof or a prodrug thereof.

27. (Withdrawn) The compound of Claim 26 wherein:

x and y are each 1;

V_a is $-C(O)-$;

R^1 is hydrogen or C_1 - C_6 alkyl;

R^2 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_1 - C_6 alkoxy, C_3 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

R^3 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_2 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl,

C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

R⁴, R⁵ and R⁶ are each hydrogen; and

R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each hydrogen.

28. (Withdrawn) The compound of Claim 27 wherein:

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆trihaloalkyl, C₁-C₆trihaloalkoxy, C₁-C₆alkylsulfonyl, -N(R¹²)₂, -OC(O)R¹², -C(O)OR¹², -S(O)₂N(R¹²)₂, cycloalkyl, heterocyclyl, heteroaryl and heteroarylcycloalkyl; and

each R¹² is independently selected from hydrogen, C₁-C₆alkyl, C₃-C₆cycloalkyl, aryl or aralkyl.

29. (Withdrawn) The compound of Claim 28 wherein:

R² is C₁-C₁₂alkyl or C₂-C₁₂alkenyl; and

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy.

30. (Withdrawn) The compound of Claim 29 selected from the group consisting of the following:

Pentane-1-sulfonic acid {5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-amide;
and

Hexane-1-sulfonic acid {5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-amide.

31. (Withdrawn) The compound of Claim 28 wherein:

R² is C₇-C₁₂aralkyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₃alkyl and C₁-C₆trihaloalkyl; and


R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy.

32. (Withdrawn) The compound of Claim 31, namely, 3-Phenyl-propane-1-sulfonic acid {5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-amide.

33. (Withdrawn) A method of treating a disease or condition mediated by stearyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 26.

34. (Withdrawn) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound of Claim 26.

35. (Withdrawn) A compound of formula (IV):



wherein:

x and y are each independently 1;

V_a is $-C(O)-$, $-C(S)-$, $-C(O)N(R^1)-$, $-C(O)O-$, $-S(O)_2-$ or $-S(O)_2N(R^1)-$;

each R^1 is independently selected from the group consisting of hydrogen, C_1 - C_{12} alkyl, C_2 - C_{12} hydroxyalkyl, C_4 - C_{12} cycloalkylalkyl and C_7 - C_{19} aralkyl;

R^2 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_3 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

or R^2 is a multi-ring structure having 2 to 4 rings wherein the rings are independently selected from the group consisting of cycloalkyl, heterocyclyl, aryl and heteroaryl, where some or all of the rings may be fused to each other;

R^3 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_2 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

or R^3 is a multi-ring structure having 2 to 4 rings wherein the rings are independently selected from the group consisting of cycloalkyl, heterocyclyl, aryl and heteroaryl and where some or all of the rings may be fused to each other;

R^4 , R^5 and R^6 are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or $-N(R^{13})_2$;

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each independently selected from hydrogen or C_1 - C_3 alkyl;

and

each R^{13} is independently selected from hydrogen or C_1 - C_6 alkyl;

a stereoisomer, enantiomer or tautomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutical composition thereof or a prodrug thereof.

36. (Withdrawn) The compound of Claim 35 wherein:

x and y are each 1;

V_a is $-C(O)-$;

each R^1 is independently hydrogen or C_1 - C_6 alkyl;

R^2 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_3 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

R^3 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl,

C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₂-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, aryl, C₇-C₁₉aralkyl, C₃-C₁₂heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

R⁴, R⁵ and R⁶ are each hydrogen; and

R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each hydrogen.

37. (Withdrawn) The compound of Claim 36 wherein:

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆trihaloalkyl, C₁-C₆trihaloalkoxy, C₁-C₆alkylsulfonyl, -N(R¹²)₂, -OC(O)R¹², -C(O)OR¹², -S(O)₂N(R¹²)₂, cycloalkyl, heterocyclyl, heteroaryl and heteroarylcycloalkyl; and

each R¹² is independently selected from hydrogen, C₁-C₆alkyl, C₃-C₆cycloalkyl, aryl or aralkyl.

38. (Withdrawn) The compound of Claim 37 wherein:

R² is C₁-C₁₂alkyl or C₂-C₁₂alkenyl; and

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy.

39. (Withdrawn) The compound of Claim 38 selected from the group consisting of the following:

1-(3-Methyl-butyl)-3-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-urea;

1-Pentyl-3-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-urea; and

1-Butyl-3-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-urea.

40. (Withdrawn) The compound of Claim 37 wherein:

R² is C₇-C₁₂aralkyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₃alkyl and C₁-C₆trihaloalkyl; and

R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl and C_1 - C_6 trihaloalkoxy.

41. (Withdrawn) The compound of Claim 40 selected from the group consisting of the following:

1-[3-(4-Fluoro-phenyl)-propyl]-3-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-urea;


1-Phenethyl-3-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-urea; and

1-Benzyl-3-{5-[4-(2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridin-2-yl}-urea.

42. (Withdrawn) A method of treating a disease or condition mediated by stearyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 35.

43. (Withdrawn) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound of Claim 35.

44. (Withdrawn) A compound of formula (V):



wherein:

x and y are each independently 1;

W_a is -O-, -N(R^1)- or -S(O)_t- (where t is 0, 1 or 2);

V_a is -C(O)-, -C(S)-, -C(O)N(R^1)-, -C(O)O-, -S(O)₂- or -S(O)₂N(R^1)-;

x and y are each independently 1, 2 or 3;

each R^1 is independently selected from the group consisting of hydrogen, C_1 - C_{12} alkyl, C_2 - C_{12} hydroxyalkyl, C_4 - C_{12} cycloalkylalkyl and C_7 - C_{19} aralkyl;

R^2 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_3 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

or R^2 is a multi-ring structure having 2 to 4 rings wherein the rings are independently selected from the group consisting of cycloalkyl, heterocyclyl, aryl and heteroaryl, where some or all of the rings may be fused to each other;

R^3 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_2 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

or R^3 is a multi-ring structure having 2 to 4 rings wherein the rings are independently selected from the group consisting of cycloalkyl, heterocyclyl, aryl and heteroaryl and where some or all of the rings may be fused to each other;

R^4 , R^5 and R^6 are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or $-N(R^{13})_2$;

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each independently selected from hydrogen or C_1 - C_3 alkyl;

and

each R^{13} is independently selected from hydrogen or C_1 - C_6 alkyl;

a stereoisomer, enantiomer or tautomer thereof, a pharmaceutically acceptable salt thereof, a pharmaceutical composition thereof or a prodrug thereof.

45. (Withdrawn) The compound of Claim 44 wherein:

x and y are each 1;

W_a is -O-;

V_a is -C(O)-;

R¹ is hydrogen or C₁-C₆alkyl;

R² is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₃-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, aryl, C₇-C₁₉aralkyl, C₃-C₁₂ heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

R³ is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₂-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, aryl, C₇-C₁₉aralkyl, C₃-C₁₂heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

R⁴, R⁵ and R⁶ are each hydrogen; and

R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each hydrogen.

46. (Withdrawn) The compound of Claim 45 wherein:

R³ is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆trihaloalkyl, C₁-C₆trihaloalkoxy, C₁-C₆alkylsulfonyl, -N(R¹²)₂, -OC(O)R¹², -C(O)OR¹², -S(O)₂N(R¹²)₂, cycloalkyl, heterocyclyl, heteroaryl and heteroarylcycloalkyl; and

each R¹² is independently selected from hydrogen, C₁-C₆alkyl, C₃-C₆cycloalkyl, aryl or aralkyl.

47. (Withdrawn) The compound of Claim 44 wherein:

x and y are each 1;

W_a is -N(R¹)-;

V_a is -C(O)-;

R^1 is hydrogen or C_1 - C_6 alkyl;

R^2 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_3 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

R^3 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_2 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

R^4 , R^5 and R^6 are each hydrogen; and

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each hydrogen.

48. (Withdrawn) The compound of Claim 47 wherein:

R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl, C_1 - C_6 trihaloalkoxy, C_1 - C_6 alkylsulfonyl, $-N(R^{12})_2$, $-OC(O)R^{12}$, $-C(O)OR^{12}$, $-S(O)_2N(R^{12})_2$, cycloalkyl, heterocyclyl, heteroaryl and heteroarylcycloalkyl; and

each R^{12} is independently selected from hydrogen, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, aryl or aralkyl.

49. (Withdrawn) The compound of Claim 44 wherein:

x and y are each 1;

W_a is $-S(O)_t$ - (where t is 0, 1 or 2);

V_a is $-C(O)-$;

R^2 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_3 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

R^3 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_2 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

R^4 , R^5 and R^6 are each hydrogen; and

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each hydrogen.

50. (Withdrawn) The compound of Claim 49 wherein:


R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl, C_1 - C_6 trihaloalkoxy, C_1 - C_6 alkylsulfonyl, $-N(R^{12})_2$, $-OC(O)R^{12}$, $-C(O)OR^{12}$, $-S(O)_2N(R^{12})_2$, cycloalkyl, heterocyclyl, heteroaryl and heteroarylalkyl; and

each R^{12} is independently selected from hydrogen, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, aryl or aralkyl.

51. (Withdrawn) A method of treating a disease or condition mediated by stearyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 44.

52. (Withdrawn) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound of Claim 44.

53. (Previously Presented) A compound of formula (VIa):



wherein:

x and y are each independently 1;

R¹ is selected from the group consisting of hydrogen, C₁-C₁₂alkyl, C₂-C₁₂hydroxyalkyl, C₄-C₁₂cycloalkylalkyl and C₇-C₁₉aralkyl;

R² is selected from the group consisting of C₇-C₁₂alkyl, C₃-C₁₂alkenyl, C₇-C₁₂hydroxyalkyl, C₂-C₁₂alkoxyalkyl, C₃-C₁₂hydroxyalkenyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, C₁₃-C₁₉aralkyl, C₃-C₁₂heterocyclalkyl, and C₃-C₁₂heteroarylalkyl;

R³ is phenyl or naphthyl;

R⁴, R⁵ and R⁶ are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or -N(R¹³)₂;

R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each independently selected from hydrogen or C₁-C₃alkyl;

and

each R¹³ is independently selected from hydrogen or C₁-C₆alkyl;

including a stereoisomer, enantiomer or tautomer thereof, or a pharmaceutically acceptable salt thereof.

54. (Previously Presented) The compound of Claim 53 wherein:

x and y are each 1;

R¹ is hydrogen or C₁-C₆alkyl;

R² is selected from the group consisting of C₇-C₁₂alkyl, C₃-C₁₂alkenyl, C₇-C₁₂hydroxyalkyl, C₂-C₁₂alkoxyalkyl, C₃-C₁₂hydroxyalkenyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, C₁₃-C₁₉aralkyl, C₃-C₁₂heterocyclalkyl, and C₃-C₁₂heteroarylalkyl;

R³ is phenyl or naphthyl;

R⁴, R⁵ and R⁶ are each hydrogen; and


R⁷, R^{7a}, R⁸, R^{8a}, R⁹, R^{9a}, R¹⁰, and R^{10a} are each hydrogen.

55. (Currently Amended) A method of treating a disease or condition mediated by

stearoyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 53, and wherein the disease or condition is selected from the group consisting of Type II diabetes, impaired glucose tolerance, insulin resistance, obesity, fatty liver, non-alcoholic steatohepatitis, dyslipidemia, acne, and ~~metabolic syndrome~~ and any combination of these.

56. (Original) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound of Claim 53.

57. (Previously Presented) A compound of formula (VIb):



wherein:

x and y are each independently 1;

each R¹ is independently selected from the group consisting of hydrogen, C₁-C₁₂alkyl, C₂-C₁₂hydroxyalkyl, C₄-C₁₂cycloalkylalkyl and C₇-C₁₉aralkyl;

R² is selected from the group consisting of C₁-C₁₂alkyl, C₂-C₁₂alkenyl, C₂-C₁₂hydroxyalkyl, C₂-C₁₂hydroxyalkenyl, C₃-C₁₂alkoxyalkyl, C₃-C₁₂cycloalkyl, C₄-C₁₂cycloalkylalkyl, aryl, C₇-C₁₉aralkyl, C₃-C₁₂heterocyclyl, C₃-C₁₂heterocyclylalkyl, C₁-C₁₂heteroaryl and C₃-C₁₂heteroarylalkyl;

R³ is naphthyl or phenyl, each optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C₁-C₆alkyl, C₁-C₆trihaloalkyl, C₁-C₆trihaloalkoxy, C₁-C₆alkylsulfonyl, -N(R¹²)₂, -OC(O)R¹², -C(O)OR¹², -S(O)₂N(R¹²)₂, cycloalkyl, heterocyclyl, heteroaryl and heteroarylalkyl, provided that R³ is not phenyl

substituted with optionally substituted thienyl, and provided that when R^3 is naphthyl, R^2 can not be C_1 - C_6 alkyl, C_2 - C_6 hydroxyalkyl or phenyl substituted by amino;

R^4 , R^5 and R^6 are each independently selected from hydrogen, fluoro, chloro, methyl, methoxy, trifluoromethyl, cyano, nitro or $-N(R^{13})_2$;

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each independently selected from hydrogen or C_1 - C_3 alkyl;

each R^{12} is independently selected from hydrogen, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, aryl or aralkyl; and

each R^{13} is independently selected from hydrogen or C_1 - C_6 alkyl;

a stereoisomer, enantiomer or tautomer thereof, or a pharmaceutically acceptable salt thereof.

58. (Original) The compound of Claim 57 wherein:

x and y are each 1;

R^1 is hydrogen or C_1 - C_6 alkyl;

R^2 is selected from the group consisting of C_1 - C_{12} alkyl, C_2 - C_{12} alkenyl, C_2 - C_{12} hydroxyalkyl, C_2 - C_{12} hydroxyalkenyl, C_3 - C_{12} alkoxyalkyl, C_3 - C_{12} cycloalkyl, C_4 - C_{12} cycloalkylalkyl, aryl, C_7 - C_{19} aralkyl, C_3 - C_{12} heterocyclyl, C_3 - C_{12} heterocyclylalkyl, C_1 - C_{12} heteroaryl and C_3 - C_{12} heteroarylalkyl;

R^3 is naphthyl or phenyl, each optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl, C_1 - C_6 trihaloalkoxy, C_1 - C_6 alkylsulfonyl, $-N(R^{12})_2$, $-OC(O)R^{12}$, $-C(O)OR^{12}$ or $-S(O)_2N(R^{12})_2$;

R^4 , R^5 and R^6 are each hydrogen;

R^7 , R^{7a} , R^8 , R^{8a} , R^9 , R^{9a} , R^{10} , and R^{10a} are each hydrogen; and

each R^{12} is independently selected from hydrogen, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, aryl or aralkyl.

59. (Original) The compound of Claim 58 wherein:

R^2 is C_7 - C_{12} alkyl optionally substituted by one or more substituents selected from the group consisting of halo, C_1 - C_3 alkyl and C_1 - C_6 trihaloalkyl; and

R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl and C_1 - C_6 trihaloalkoxy.

60. (Original) The compound of Claim 59 selected from the group consisting of the following:

5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid (3-phenyl-propyl)-amide;

5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid phenethyl-amide;

5-[4-(2-Trifluoromethylbenzoyl)piperazin-1-yl]pyridine-2-carboxylic acid [2-(4-fluorophenyl)ethyl]amide;

5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid [3-(4-fluorophenyl)-propyl]-amide;

5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid 4-trifluoromethylbenzylamide;

5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid [3-(4-trifluoromethyl-phenyl)-propyl]-amide; and

5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid [2-(4-trifluoromethyl-phenyl)-ethyl]-amide.

61. (Original) The compound of Claim 58 wherein:

R^2 is C_1 - C_{12} alkyl or C_2 - C_{12} alkenyl; and

R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl and C_1 - C_6 trihaloalkoxy.

62. (Original) The compound of Claim 61 selected from the group consisting of the following:

5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid (3-methyl-butyl)-amide;
5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid hexylamide;
5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid pentylamide;
5-[4-(4-Fluoro-2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid (3-methyl-butyl)-amide; and
5-[4-(5-Fluoro-2-trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid (3-methyl-butyl)-amide.

63. (Original) The compound of Claim 58 wherein:
 R^2 is C_3 - C_{12} cycloalkyl or C_4 - C_{12} cycloalkylalkyl; and
 R^3 is phenyl optionally substituted by one or more substituents selected from the group consisting of halo, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl and C_1 - C_6 trihaloalkoxy.

64. (Original) The compound of Claim 63 selected from the group consisting of the following:
5-[4-(2-Trifluoromethylbenzoyl)piperazin-1-yl]pyridine-2-carboxylic acid (3-cyclohexyl-propyl)amide;
5-[4-(6-Trifluoromethyl-cyclohexa-1,3-dienecarbonyl)-piperazin-1-yl]-pyridine-2-carboxylic acid (2-cyclohexyl-ethyl)-amide; and
5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carboxylic acid cyclohexylmethyl-amide.

65. (Original) The compound of Claim 58 wherein:
 R^2 is C_3 - C_{12} heterocyclylalkyl optionally substituted by one or more substituents selected from the group consisting of halo, cyano, nitro, hydroxy, C_1 - C_6 alkyl, C_1 - C_6 trihaloalkyl, C_1 - C_6 trihaloalkoxy, C_1 - C_6 alkylsulfonyl, $-N(R^{12})_2$, $-OC(O)R^{12}$, $-C(O)OR^{12}$ and $-S(O)_2N(R^{12})_2$;
 R^3 is phenyl optionally substituted by one or more substituents selected from the

group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy; and
each R¹² is independently selected from hydrogen, C₁-C₆alkyl, C₃-C₆cycloalkyl, aryl or aralkyl.

66. (Original) The compound of Claim 65 wherein R² is 2-piperazinylethyl optionally substituted by -C(O)OR¹².

67. (Original) The compound of Claim 66, namely, 4-[2-({5-[4-(2-Trifluoromethyl-benzoyl)-piperazin-1-yl]-pyridine-2-carbonyl}-amino)-ethyl]-piperazine-1-carboxylic acid tert-butyl ester.

68. (Original) The compound of Claim 58 wherein:
R² is C₇-C₁₂aralkyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₃alkyl and C₁-C₆trihaloalkyl; and
R³ is naphthyl optionally substituted by one or more substituents selected from the group consisting of halo, C₁-C₆alkyl, C₁-C₆trihaloalkyl and C₁-C₆trihaloalkoxy.

69. (Original) The compound of Claim 68 selected from the group consisting of the following:

5-[4-(Naphthalene-1-carbonyl)-piperazin-1-yl]-pyridine-2-carboxylic acid (3-phenyl-propyl)-amide; and

5-[4-(Naphthalene-1-carbonyl)piperazin-1-yl]pyridine-2-carboxylic acid phenethylamide.

70. (Currently Amended) A method of treating a disease or condition mediated by stearoyl-CoA desaturase (SCD) in a mammal, wherein the method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 57, wherein the disease or condition is selected from the group consisting of Type II diabetes, impaired glucose tolerance, insulin resistance, obesity, fatty liver, non-alcoholic steatohepatitis,

dyslipidemia, acne, and ~~metabolic syndrome~~ and any combination of these.

71. (Original) A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of a compound of Claim 57.